

A rare neuroendocrine tumor of the lung: sclerosing paraganglioma. A neoplasm that is difficult to diagnose and a source of dangerous pitfalls. A case report and literature review

Giovanni Africa¹, Francesca M. Plutino¹, Marcello Filotico²

¹ Grande Ospedale Metropolitano "Bianchi-Melacrino-Morelli, Reggio Calabria, Italy; ² Fondazione Cad. Panico, Tricase, LE, Italy

Summary

An endobronchial obstructing neof ormation was found in a 58-year-old man. The histology and immunohistochemical profile oriented the authors towards a diagnosis of paraganglioma, sclerosing variant. This very difficult diagnosis, especially in a pulmonary localization, may lead to erroneous conclusions both in terms of histogenetic interpretation and that of its biological behavior. The pulmonary localization of the paraganglioma is very rare and even more rare the sclerosing variant, recently reported. Differential diagnosis and literature are discussed.

Key words: paraganglioma, immunohistochemistry

Introduction

Paraganglioma (PG) is the extra adrenal analogue of pheochromocytoma of which it repeats the structure, immunocytochemical profile and biological behavior. It is a rare neoplasm with various localizations among which the pulmonary one is very rare. The case we observed, beyond the pulmonary site, presented the histomorphological features of the very rare sclerosing variant. It is a neoplasia of very difficult diagnosis prone to dangerous pitfalls. For all these reasons, we considered it worthy of reporting.

Case – In a man of 58 years, with ischemic heart disease, bearer of a pacemaker, to a radiological control with CT Scan, the presence of a neof ormation vegeto infiltrating obstructing the anterior segmental bronchus of the upper lobe of the right lung was detected (Figs. 1a,b). Bronchoscopic examination revealed the presence of a large base polypoid neof ormation covered by a glossy mucosa, brownish in color almost entirely occupying the bronchial lumen (Fig. 2a). A large bioptic resection was performed on the neof ormation.

After biopsy, (July 2019) and histopathologic diagnosis, the patient no longer scheduled for subsequent treatment and follow-up.

Material and methods

The biopsy specimen was fixed in formalin and embedded in paraffin. The

Received: August 30, 2019
Accepted: September 17, 2020

Correspondence

Marcello Filotico
Fondazione Cad. Panico, Tricase, LE, Italy
E-mail: mfilotico@gmail.com

Conflict of interest

The Authors declare no conflict of interest.

How to cite this article: Africa G, Plutino FM, Filotico M. A rare neuroendocrine tumor of the lung: sclerosing paraganglioma. A neoplasm that is difficult to diagnose and a source of dangerous pitfalls. A case report and literature review. *Pathologica* 2020;112:191-194. <https://doi.org/10.32074/1591-951X-23-19>

© Copyright by Società Italiana di Anatomia Patologica e Citopatologia Diagnostica, Divisione Italiana della International Academy of Pathology



OPEN ACCESS

This is an open access journal distributed in accordance with the CC-BY-NC-ND (Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International) license: the work can be used by mentioning the author and the license, but only for non-commercial purposes and only in the original version. For further information: <https://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>




Figure 1. (a-b) Axial Total body TC scan with contrast medium. The truncation of the right upper lobar bronchus is evident.

Table I. Immunohistochemical panel.

Ab	Dilutions
Vimentin	Monoclonal 1:50
Cytokeratin MNF 116	Monoclonal 1:50
S100	Polyclonal 1:400
Chromogranin A	Monoclonal 1:100
Synaptophysin	Monoclonal 1:50
NSE	Monoclonal 1:100
CD56	Monoclonal 1:50
TTF1	Monoclonal 1:100
Ki67	Monoclonal 1:75
CD34	Monoclonal 1:20
CD31	Monoclonal 1:20
p40	Monoclonal 1:100
SMACT	Monoclonal 1:50
GFAP	Polyclonal 1:200
PD-1	Monoclonal 1:50
CD117	Polyclonal 1:400




Figure 2. (a) Bronchoscopic image. Polypoid neformation occluding the bronchial lumen; (b) Dome-shaped surgical specimen; (c-d). Subtiles cell chains compressed by exuberant desmoplasia HE 175X.




Figure 3. (a-b) Branched cellular cords surrounded by an abundant collagen matrix. HE 150, 175 X; (c-d) - *Zell Ballen*. HE 150, 175 X.

sections were stained with hematoxylin-eosin, Masson's trichrome, and Silver-impregnation according to Gomori. For immunohistochemistry a panel of antibodies was used (Tab. I).

Histology

The material under examination consisted of a grossly polypoid fragment with a maximum diameter of 4 mm (Fig. 2b). The mass was totally ulcerated. Adjacent were a few minute frustule of bronchial mucosa. The

tissue was mainly made up of a proliferation of medium-sized globose elements, voluminous roundish hyperchromatic nucleus, mainly arranged in chains (Figs. 2c,d) or joined to form thin branched cords compressed and separated by a dense and abundant fibrous matrix (Figs. 3a,b). In the periphery, where the fibrous tissue was more lax and less abundant, elements aggregated into nodular formations type *Zell Ballen* (Figs. 3c,d). Silver staining according to Gomori allows recognition, although distorted, the characteristic alveolar pattern of the PG (Figs. 4a), while Masson's trichrome confirmed the intense desmoplasia (Figs. 4b). Mitotic activity was practically absent.

The results of the immunohistochemical investigation are shown in Table II.

On the basis of morphological data and the results of immunohistochemical investigation, after having examined a vast differential diagnosis panel, a diagnosis of primary broncho-pulmonary paraganglioma, sclerosing variant was favored.

Discussion

The adrenal medulla and the extra adrenal paraganglia, having the same embryological origin, the same histological structure and the same function constitute the adrenal sympathetic neuroendocrine system. The extra adrenal paraganglia are structures distributed along the paraortic-praventebral axis, in parallel with the distribution of the sympathetic nervous system. They produce neurotransmitters (epinephrine and norepinephrine) that activate the interneuronal synapses of the sympathetic system. The paraganglia with respect to the site are divided into branchiomic, intravagal, aortosympatetic and visceral. The common origin from the Neural crest with the cells of the diffuse neuroendocrine system explains the common dyeing, histochemi-

cal, immunohistochemical and ultrastructural affinity, making it, especially in lung localizations, very difficult for diagnostic differentiation¹.

Pulmonary localization of PG is among the rarest. In a collection of 152 cases, the pulmonary site appears only in 3 (2%)². A review of the literature in 1995 reported 25 cases up to that date³. Subsequent to that time another 13 have been reported in the literature in the English language³⁻¹⁴.

The differential diagnosis of this lesion in the lung with more frequent carcinoid tumors is not always easy and lies in subtle morphological differences and immunophenotypic expressiveness. So much so that according to some AA, while not excluding the possibility of their existence in such a location, it is very difficult to determine their actual frequency given the difficulty differential diagnosis with carcinoid¹. On the morphological level differential diagnosis can be fairly easy in the case of lesions in the classical form. It becomes very difficult if not impossible in the atypical or undifferentiated forms or in the case of not uncommon variants.

On the basis of immunophenotypic differentiation it is a little less arduous as can be seen from Table III.

Although previous cases of PG with marked sclerosis phenomena have been reported, a specific sclerosing variant was defined in 2006¹⁵. In that publication, the result of a multicentric study conducted at four large North American institutions, 16 cases were collected and described. Subsequent to this first publication, five other individual cases followed, one of which was bronchopulmonary¹⁶⁻²⁰.

The morphological patterns presented in these publications are completely superimposable to the one observed to us. Our differential diagnostic procedure followed the same scheme proposed in these articles. In view of the lack of the classic PG structure, the infiltrative aspect of the lesion, one can consider the possibility of a primitive or secondary pulmonary malignancy, and among these those with desmoplasia. Secondly, the neuroendocrine expression introduces a discussion between the possible neuroendocrine lesions of the lung. The morphological picture as a whole does not integrate the typical aspects of any of the neuroendocrine neoplasms of the lung. In addition, in the peripheral areas, less affected by desmoplasia, the characteristic *Zell Ballen* of the PG are easily recognizable (Figs. 3c-




Figure 4. (a) Silver staining - Despite the thickening of the septa, the characteristic alveolar structure of the PG is still recognizable 150, 250 X; (b) Masson Trichrome intense desmoplasia 175 X; (c) Chromogranin; (d) Synaptophysin - 200X.

Table II. Immunohistochemistry results.

Ab	Vim	Ck	S100	Chromogr	Synapto	NSE	CD56	TTF1	Ki67	CD34	CD31	P40	SMACT	GFAP	PD-1	CD117
	+	-	+	+	+	+	+/-	-	low	-	-	-	-	-	-	-
Fig.	5d		5c	4c	4d	5a	5b									

*sustentacular cells.

Table III. PG vs carcinoid immunohistochemistry.

Ab	Parganglioma	CARCINOID
CK	.	+
SYNAPTO	+	+
CHROMOGR	+	+
NSE	+	+
TTF1	-	+
S100	+*	-

**sustentacular cells*.




Figure 5. (a) NSE; (b) CD56; (c) S100 (sustentacular cells) 150X; (d) Vimentin 200X.

d). In addition, the silver staining highlights, albeit distorted, the characteristic alveolar pattern (Fig. 4a). On the immunohistochemical level, despite the clearly positive neuroendocrine markers (Figs. 4c,d, 5a,b), the negativity for cytokeratins and TTF1 makes a neoplasm of the carcinoid family unlikely. Even if scattered, because of sclerosis, but still placed on the periphery of the cellular cords, positive S100 cells are present interpretable as *sustentacular* (Fig. 5c). Vimentin has an arrangement similar to that of the argyrophilic fibers that surround the alveolar spaces. The low proliferation index and the absence of mitotic figures suggest that the lesion is low in aggressiveness. The incomplete removal, if not followed by a radicalizing intervention, could cause recurrence.

References

- Tumors of Adrenal Glands and Extraadrenal Paraganglia. AFIP Atlas of Tumor Pathology, Series 4. Washington DC: ARP 2007, p. 283.
- Feng N, Zhang WY, Wu XT. Clinicopathological analysis of paraganglioma with literature review. *World J Gastroenterol* 2009;15:3003-8. <https://doi.org/10.3748/wjg.15.3003>
- Skodt V, Jacobsen GK, Helsted M. Primary paraganglioma of the lung: report of two cases and review of literature. *APMIS* 1995;103:597-603. <https://doi.org/10.1111/j.1699-0463.1995.tb01412.x>
- Hironaka M, Fukayama M, Takayashiki N, et al. Pulmonary gangliocytic paraganglioma: case report and comparative immunohistochemical study of related neuroendocrine neoplasms. *Am J Surg Pathol* 2001;25:688-93. <https://doi.org/10.1097/0000478-200105000-00020>
- da Silva RA, Gross JL, Haddad FJ, et al. Primary pulmonary paraganglioma: case report and literature review. *Clinics (Sao Paulo)* 2006;61:83-6. <https://doi.org/10.1590/s1807-59322006000100015>
- Kim KN, Lee KN, Roh MS, Choi PJ, et al. Pulmonary paraganglioma manifesting as an endobronchial mass. *Korean J Radiol* 2008;9:87-90. <https://doi.org/10.3348/kjr.2008.9.1.87>
- Ceberut K, Müslehiddinoglu A, Akar I, et al. Primary pulmonary paraganglioma: a case report. *Turk Jem* 2010;14:14-6.
- Huwer H, Kalweit G, Schäfer H, et al. Pulmonary paraganglioma: case report and literature review. *Pneumologie* 2011;65:742-4. <https://doi.org/10.1055/s-0031-1286635>
- Zhang JJ, Liu T, Peng F. Primary paraganglioma of the lung: a case report and literature review. *J Int Med Res* 2012;40:1617-26. <https://doi.org/10.1177/147323001204000442>
- Ibraheema T, ElGhazaly H, Madkoura A, et al. Primary paraganglioma of the lung: a case report. *Egyptian Journal of Bronchology* 2014 8:64-5.
- Huang X, Liang QL, Jiang Let al. Primary pulmonary paraganglioma: a case report and review of literature. *Medicine* 2015;94:e1271. <https://doi.org/10.1097/MD.0000000000001271>
- Fiorentino G, Annunziata A, De Rosa N. Primary paraganglioma of the lung: a case report. *J Med Case Rep* 2015;9:166. <https://doi.org/10.1186/s13256-015-0639-z>
- Chan J, Sangani R, Oduntan O, et al. Primary Pulmonary paraganglioma presenting as a solitary lung nodule. *150#4S Chest* october 2016.
- Muriana P, Bandiera A, Ciriaco Pet al. A case of endobronchial paraganglioma. *Ann R Coll Surg Engl* 2017;99:e28-e30. <https://doi.org/10.1308/rcsann.2016.0284>
- Plaza JA, Wakely PE Jr, Moran Cet al. Sclerosing paraganglioma: report of 19 cases of an unusual variant of neuroendocrine tumor that may be mistaken for an aggressive malignant neoplasm. *Am J Surg Pathol* 2006;30:7-12. <https://doi.org/10.1097/01.pas.0000174012.37439.c7>
- Mukhopadhyay S, Naqvi AH, Su W, et al. Airway-occlusive sclerosing paraganglioma presenting with dyspnea. *Am J Surg Pathol* 2006;30:1206-7. <https://doi.org/10.1097/01.pas.0000209844.67293.88>
- Evankovich J, Dedhia RC, Bastaki JM, et al. Primary sclerosing paraganglioma of the thyroid gland: a case report. *Ann Otol Rhinol Laryngol* 2012;121:510-5. <https://doi.org/10.1177/000348941212100803>
- Santi R, Franchi A, Saladino V, et al. Sclerosing paraganglioma of the carotid body: a potential pitfall of malignancy. *Head Neck Pathol* 2015;9:300-4. <https://doi.org/10.1007/s12105-014-0569-x>
- Ng E, Duncan G, Choong AM, et al. Sclerosing paragangliomas of the carotid body: a series of a rare variant and review of the literature. *Ann Vasc Surg* 2015(7):1454.e5-1454.e12. <https://doi.org/10.1016/j.avsg.2015.04.083>
- Blefaria NDA, Eliezer DD, Vilain RE, et al. Sclerosing paraganglioma of the carotid body in a child: A diagnostic and therapeutic challenge. *International Journal of Pediatric Otorhinolaryngology Case Reports* 2019;23:100663. <https://doi.org/10.1016/j.pedeo.2019.100663>